



Stony Brook
Medicine

Colonoscopy Preparation Instructions: Two Day SUPREP Bowel Prep

Name: _____

Procedure: Colonoscopy _____

Physician: _____

Location: _____

Date of Procedure: _____

Arrive at: _____

Please note: As a courtesy to other patients and to your doctor, please call (631) 444-5220 as soon as possible if you are unable to keep your appointment.



AS SOON AS POSSIBLE

1. Make arrangements for a responsible adult to accompany you when leaving the procedure, even if taking public transportation. **Your procedure may be canceled if we cannot confirm your transportation on the day of the procedure. Taxi and Rideshare (Uber®, Lyft® etc.) drivers are not acceptable.**

******IMPORTANT: IF YOU DO NOT HOLD CERTAIN MEDICATION(S) AS INSTRUCTED BELOW OR OBTAIN ANY NECESSARY CLEARANCES, YOUR PROCEDURE MAY BE CANCELED. IT IS YOUR RESPONSIBILITY THAT CLEARANCES ARE FAXED TO 631-444-5225 WITH DR.'S NAME AND DATE OF PROCEDURE**

2. If you take any medications that increase your risk of bleeding, you must ask the prescribing doctor when to stop taking the medication before the procedure. The most common of these medications include but are not limited to:

Warfarin (Coumadin®)	Effient (Prasugrel®)
Clopidogrel (Plavix®)	Apixaban (Eliquis®)
Ticagrelor (Brilinta®)	Dabigatran (Pradaxa®)
Rivaroxaban (Xarelto®)	Enoxaparin (Lovenox®)

You may continue taking aspirin unless otherwise instructed

3. For patients with diabetes
 - If you take insulin, contact your diabetes doctor a minimum of **14 days prior to your procedure date** to adjust your insulin dose during the bowel preparation
 - If you use an insulin pump, a **written letter** from your endocrinologist giving clear instructions on adjustments for basal rate before and during the procedure will be required (**the letter must be obtained at least 14 days prior to your procedure date**)
 - If you use oral diabetes medications (except an SGLT-2 inhibitor), do not take these medications on the morning of your procedure; you may resume this medication once you resume a regular diet.
 - If you take an **SGLT-2 inhibitor** (Canigliflozin [Invokana], Dapagliflozin [Farxiga], or Empagliflozin [Jardiance]) **for diabetes**, do not take this medication for **3 days prior to your procedure date**. If you take Ertugliflozin [Steglatro], do not take this medication for **4 days prior to your procedure date**. After your procedure, you may take this medication once you resume a regular diet. If this medication is prescribed for a **cardiac indication**, please discuss holding this medication with your Cardiologist.



4. For patients taking a GLP-1 Agonist for diabetes or obesity

Drug	Interruption of Therapy	Resumption of Therapy
Once Daily GLP-1RA (SQ) Liraglutide (Victoza®, Saxenda®)	Hold day of procedure	Resume next day
Twice-Daily GLP-1RA (SQ) Exenatide (Byetta®)	Hold the night before procedure and the morning of the procedure	Resume next day/next dose
Once Daily GLP-1RA (Oral) Semaglutide (Rybelsus®)	Hold one week before the procedure	Resume next day
Weekly GLP-1 RA (SQ) Dulaglutide (Trulicity®) Exenatide (Bydureon ®) Semaglutide (Ozempic®, Wegovy®) Tirzapatide (Mounjaro®)	Hold one week before the procedure	Resume next day /next dose

5. If you have a pacemaker or defibrillator, please make sure your doctor is aware and bring the device card with you to your procedure



5 DAYS PRIOR TO COLONOSCOPY

- Buy one 119g bottle of Miralax®
- Pick up your prescribed SUPREP bowel prep kit from the pharmacy
- Go to www.suprepkit.com/voucher.htm for a savings coupon
- Stop taking any iron supplements, herbal medications, or fish oil
- Open the 119g bottle of Miralax and take 1 capful dissolved in water, juice or coffee once a day until day before colonoscopy
- Avoid eating any nuts or seeds including poppy seeds, sesame seeds and sunflower seeds, any fruits or vegetables with seeds, or corn.
- Call the Admitting Office at (631) 444-1870 and provide your insurance information
- Confirm your transportation for the day of your procedure
- Please call if you need to reschedule your procedure

2 DAYS PRIOR TO COLONOSCOPY

- **Do not eat any solid or soft food. Drink plenty of clear liquids throughout the day to prevent dehydration from the bowel preparation. Do not drink alcohol.**

NOTE: If you have Type 1 diabetes, please discuss an acceptable clear liquid diet with your Endocrinologist.

Examples of clear liquids (able to be seen through when poured into a glass) include:

- | | |
|--|---|
| ✓ Water | ✓ Gatorade®, Powerade® or similar sports drink (no red or purple) |
| ✓ Clear juices without pulp (apple or white grape juice) | ✓ Ginger-ale or Lemon-lime soda (7-Up® or Sprite®) |
| ✓ Clear soup broth or bouillon | ✓ Jell-O® (no red or purple) |
| ✓ Tea or coffee (1 cup) | ✓ Popsicles (no red or purple) |

- **Take 17 g (1 capful) of Miralax 3 times mixed in a clear liquid on this day (1 cap in the morning, 1 cap in the afternoon, and 1 cap in the evening) – you may use the 119g Miralax bottle for this portion of the bowel preparation.**

1 DAY BEFORE YOUR COLONOSCOPY

- Continue the same liquid diet you took the day prior
- **At 6PM** pour 1 bottle of SUPREP into the mixing container and add cool drinking water to the 16-ounce line on the container and mix. Drink ALL the liquid in the container. Drink 2 more 16-ounce containers of water over the next hour
 - ✓ It is normal to feel full and bloated during the prep
 - ✓ If you experience vomiting, stop taking the prep, wait 30 minutes and then restart and drink it slowly. If you vomit again, call our office (631) 444-5220
- Continue to drink clear liquids throughout the evening



ON THE DAY OF YOUR COLONOSCOPY

Before leaving home

- **6 hours before your scheduled procedure time**, take the second dose of SUPREP. Pour the second bottle of SUPREP into the mixing container and add cool drinking water to the 16-ounce line on the container and mix. Drink ALL the liquid in the container. Drink two additional 16-ounce containers of water over the next hour. Depending on the time of your procedure, this might be in the middle of the night. Correct timing is essential for an optimal bowel prep – set an alarm clock if necessary.
- **All prep and clear liquids should be finished 5 hours before the procedure time**
- **No gum or hard candies**
- **Take your heart and blood pressure medications with a sip of water on the day of your procedure.**
- **Do not take any oral diabetes medications; discuss insulin dosing with your Endocrinologist or Primary Care Physician**
- Bring a list of your current medications
- Bring your defibrillator/pacemaker card if you have one
- Wear comfortable loose fitting clothing and flat shoes or sneakers.
- Do not wear jewelry or bring any valuables
- Do not wear contact lenses

FOR PROCEDURES AT THE HOSPITAL

- Park in the main visitor parking garage; validation will be provided for one vehicle. Valet parking **IS NOT** eligible for validation. **You must bring the parking ticket to endoscopy for validation.**
- Report to registration on hospital level 5 (lobby) 1 hour before your scheduled procedure time. They will then send you to endoscopy on 14 North.

FOR PROCEDURES AT THE AMBULATORY CARE PAVILION (ACP)

- Directions and map to the ACP building can be found on the next page
- You can park at the gated parking lot in front of the building. **You must bring the parking ticket to endoscopy for validation.**
- **Check in at the front desk 1 hour before your procedure time and they will direct you to endoscopy**

You should expect to be at the hospital or ACP for approximately 3 hours from time of registration to discharge from the endoscopy suite

**The Ambulatory Care Pavilion at Stony Brook Medicine.
3 Edmund Pellegrino Road, Stony Brook NY**

Directions:

- Heading north on Nicolls Road, turn right onto E Loop Road. Then, make a left at the first traffic light onto Edmund D Pellegrino Road.
- While on Edmund D Pellegrino Road, head towards the signs that say “Breast Center” (we are located in the same building as the Breast Center/Old Cancer Center).
- Drive through the 1st traffic circle and continue straight.
- At the 2nd traffic circle, make a U-turn so you are going the opposite direction (see attached map for details).
- Veer to the right and you will see an entrance to a small gated parking lot in front of the building.
- Please press the button and take a parking ticket that we will validate for you. If the gate is up, drive through it and park.
- The front desk will ask you for your name and will direct you to endoscopy

